

NUCLEOTIDE SEQUENCES CODING FOR THE ppgK GENE

[0001] BACKGROUND OF THE INVENTION

The invention provides nucleotide sequences from coryneform bacteria which code for the ppgK gene and a process for the fermentative preparation of amino acids using bacteria in which the endogenous ppgK gene is enhanced. All references cited herein are expressly incorporated by reference. Incorporation by reference is also designated by the term "I.B.R." following any citation.

[0002] L-Amino acids, in particular L-lysine, are used in human medicine and in the pharmaceuticals industry, in the foodstuffs industry and very particularly in animal nutrition.

[0003] It is known that amino acids are prepared by fermentation from strains of coryneform bacteria, in particular *Corynebacterium glutamicum*. Because of their great importance, work is constantly being undertaken to improve the preparation processes. Improvements to the process can relate to fermentation measures, such as, for example, stirring and supply of oxygen, or the composition of the nutrient media, such as, for example, the sugar concentration during the fermentation, or the working up to the product form by, for example, ion exchange chromatography, or the intrinsic output properties of the microorganism itself.

[0004] Methods of mutagenesis, selection and mutant selection are used to improve the output properties of these microorganisms. Strains which are resistant to antimetabolites or are auxotrophic for metabolites of regulatory importance and produce amino acids are obtained in this manner.

[0005] Methods of the recombinant DNA technique have also been employed for some years for improving the strain of Corynebacterium strains which produce L-amino acid, by amplifying individual amino acid biosynthesis genes and investigating the effect on the amino acid production.

[0006] The invention provides new measures for improved fermentative preparation of amino acids.

[0007] BRIEF SUMMARY OF THE INVENTION

Where L-amino acids or amino acids are mentioned in the following, this means one or more amino acids, including their salts, chosen from the group consisting of L-asparagine, L-threonine, L-serine, L-glutamate, L-glycine, L-alanine, L-cysteine, L-valine, L-methionine, L-isoleucine, L-leucine, L-tyrosine, L-phenylalanine, L-histidine, L-lysine, L-tryptophan and L-arginine. L-Lysine is particularly preferred.

[0008] When L-lysine or lysine are mentioned in the following, not only the bases but also the salts, such as e.g. lysine monohydrochloride or lysine sulfate, are meant by this.

[0009] The invention provides an isolated polynucleotide from coryneform bacteria, comprising a polynucleotide sequence which codes for the ppgK gene, chosen from the group consisting of

- a) polynucleotide which is identical to the extent of at least 70% to a polynucleotide which codes for a polypeptide which comprises the amino acid sequence of SEQ ID No. 2,
- b) polynucleotide which codes for a polypeptide which comprises an amino acid sequence which is identical to the extent of at least 70% to the amino acid sequence of SEQ ID No. 2,

- c) polynucleotide which is complementary to the polynucleotides of a) or b), and
- d) polynucleotide comprising at least 15 successive nucleotides of the polynucleotide sequence of a), b) or c),

the polypeptide preferably having the activity of polyphosphate glucokinase.

[0010] The invention also provides the above-mentioned polynucleotide, this preferably being a DNA which is capable of replication, comprising:

- (i) the nucleotide sequence shown in SEQ ID No. 1, or
- (ii) at least one sequence which corresponds to sequence (i) within the range of the degeneration of the genetic code, or
- (iii) at least one sequence which hybridizes with the sequence complementary to sequence (i) or (ii), and optionally
- (iv) sense mutations of neutral function in (i).

[0011] The invention also provides

- a polynucleotide, in particular DNA, which is capable of replication and comprises the nucleotide sequence as shown in SEQ ID No. 1;
- a polynucleotide which codes for a polypeptide which comprises the amino acid sequence as shown in SEQ ID No. 2;
- a vector containing the polynucleotide according to the invention, in particular a shuttle vector or plasmid vector, and

coryneform bacteria which contain the vector or in which the endogenous ppgK gene is enhanced.

[0012] The invention also provides polynucleotides which substantially comprise a polynucleotide sequence, which are obtainable by screening by means of hybridization of a corresponding gene library of a coryneform bacterium, which comprises the complete gene or parts thereof, with a probe which comprises the sequence of the polynucleotide according to the invention according to SEQ ID No.1 or a fragment thereof, and isolation of the polynucleotide sequence mentioned.

[0013] DETAILED DESCRIPTION OF THE INVENTION

Polynucleotides which comprise the sequences according to the invention are suitable as hybridization probes for RNA, cDNA and DNA, in order to isolate, in the full length, nucleic acids or polynucleotides or genes which code for polyphosphate glucokinase or to isolate those nucleic acids or polynucleotides or genes which have a high similarity of sequence with that of the ppgK gene. They are also suitable for incorporation in so-called "arrays", "micro arrays" or "DNA chips" in order to detect and determine the corresponding polynucleotides.

[0014] Polynucleotides which comprise the sequences according to the invention are furthermore suitable as primers with the aid of which DNA of genes which code for polyphosphate glucokinase can be prepared by the polymerase chain reaction (PCR).

[0015] Such oligonucleotides serving as probes or primers contain at least 25, 26, 27, 28, 29 or 30, preferably at least 20, 21, 22, 23 or 24, and most particularly preferably at least 15, 16, 17, 18 or 19 successive nucleotides. Also suitable are oligonucleotides with a length of at least 31, 32, 33, 34, 35, 36, 37, 38, 39 or 40, or at least 41, 42, 43, 44, 45, 46, 47, 48, 49 or 50

nucleotides. Also optionally suitable are oligonucleotides with a length of at least 100, 150, 200, 250 or 300 nucleotides.

[0016] "Isolated" means separated out of its natural environment.

[0017] "Polynucleotide" in general relates to polyribonucleotides and polydeoxyribonucleotides, it being possible for these to be non-modified RNA or DNA or modified RNA or DNA.

[0018] The polynucleotides according to the invention include a polynucleotide according to SEQ ID No. 1 or a fragment produced therefrom, and also polynucleotides that are at least 70% to 80%, preferably at least 81% to 85%, particularly preferably at least 86% to 90%, and most particularly preferably at least 91%, 93%, 95%, 97% or 99% identical to the polynucleotide according to SEQ ID No. 1 or a fragment produced therefrom.

[0019] "Polypeptides" are understood as meaning peptides or proteins which comprise two or more amino acids bonded via peptide bonds.

[0020] The polypeptides according to the invention include a polypeptide according to SEQ ID No. 2, in particular those with the biological activity of polyphosphate glucokinase, and also those that are at least 70% to 80%, preferably at least 81% to 85%, particularly preferably at least 86% to 90%, and most particularly preferably at least 91%, 93%, 95%, 97% or 99% identical to the polypeptide according to SEQ ID No. 2 and that have the aforementioned activity.

[0021] The invention furthermore relates to a process for the fermentative preparation of amino acids chosen from the group consisting of L-asparagine, L-threonine, L-serine, L-glutamate, L-glycine, L-alanine, L-cysteine, L-valine, L-

methionine, L-isoleucine, L-leucine, L-tyrosine, L-phenylalanine, L-histidine, L-lysine, L-tryptophan and L-arginine using coryneform bacteria which in particular already produce amino acids and in which the nucleotide sequences which code for the ppgK gene are enhanced, in particular over-expressed.

[0022] The term "enhancement" describes in this connection the raising of the intracellular activity of one or more enzymes (proteins) in a microorganism that are coded by the corresponding DNA, by for example increasing the number of copies of the gene or genes, using a strong promoter, or using a gene or allele that codes for a corresponding enzyme (protein) having a high activity, and optionally combining these measures.

[0023] By such enhancement measures, in particular overexpression, the activity or concentration of the corresponding protein is in general raised by at least 10%, 25%, 50%, 75%, 100%, 150%, 200%, 300%, 400% or 500%, at most up to 1000% or 2000%, referred to that of the wild type protein and/or to the activity or concentration of the protein in the starting microorganism.

[0024] The microorganisms which the present invention provides can produce L-amino acids from glucose, sucrose, lactose, fructose, maltose, molasses, starch, cellulose or from glycerol and ethanol. They can be representatives of coryneform bacteria, in particular of the genus Corynebacterium. Of the genus Corynebacterium, there may be mentioned in particular the species Corynebacterium glutamicum, which is known among experts for its ability to produce L-amino acids.

[0025] Suitable strains of the genus Corynebacterium, in particular of the species Corynebacterium glutamicum (C. glutamicum), are in particular the known wild-type strains

Corynebacterium glutamicum ATCC13032
 Corynebacterium acetoglutamicum ATCC15806
 Corynebacterium acetoacidophilum ATCC13870
 Corynebacterium thermoaminogenes FERM BP-1539
 Corynebacterium melassecola ATCC17965
 Brevibacterium flavum ATCC14067
 Brevibacterium lactofermentum ATCC13869 and
 Brevibacterium divaricatum ATCC14020

and L-amino acid-producing mutants or strains prepared therefrom.

[0026] The new ppgK gene from *C. glutamicum* which codes for the enzyme polyphosphate glucokinase (EC 2.7.1.63) has been isolated.

[0027] To isolate the ppgK gene or also other genes of *C. glutamicum*, a gene library of this microorganism is first set up in *Escherichia coli* (*E. coli*). The setting up of gene libraries is described in generally known textbooks and handbooks. The textbook by Winnacker: *Gene und Klone, Eine Einführung in die Gentechnologie* [Genes and Clones, An Introduction to Genetic Engineering] (Verlag Chemie, Weinheim, Germany, 1990) *I.B.R.*, or the handbook by Sambrook et al.: *Molecular Cloning, A Laboratory Manual* (Cold Spring Harbor Laboratory Press, 1989) *I.B.R.* may be mentioned as an example. A well-known gene library is that of the *E. coli* K-12 strain W3110 set up in λ vectors by Kohara et al. (*Cell* 50, 495-508 (1987)) *I.B.R.* Bathe et al. (*Molecular and General Genetics*, 252:255-265, 1996) *I.B.R.* describe a gene library of *C. glutamicum* ATCC13032, which was set up with the aid of the cosmid vector SuperCos I (Wahl et al., 1987, *Proceedings of the National Academy of Sciences USA*, 84:2160-2164 *I.B.R.*) in the *E. coli* K-12 strain NM554 (Raleigh et al., 1988, *Nucleic Acids Research* 16:1563-1575 *I.B.R.*).

[0028] Börmann et al. (Molecular Microbiology 6(3), 317-326) (1992)) *I.B.R.* in turn describe a gene library of *C. glutamicum* ATCC13032 using the cosmid pH79 (Hohn and Collins, *Gene* 11, 291-298 (1980) *I.B.R.*).

[0029] To prepare a gene library of *C. glutamicum* in *E. coli* it is also possible to use plasmids such as pBR322 (Bolivar, *Life Sciences*, 25, 807-818 (1979) *I.B.R.*) or pUC9 (Vieira et al., 1982, *Gene*, 19:259-268 *I.B.R.*). Suitable hosts are, in particular, those *E. coli* strains which are restriction- and recombination-defective. An example of these is the strain DH5 α mc^r, which has been described by Grant et al. (Proceedings of the National Academy of Sciences USA, 87 (1990) 4645-4649) *I.B.R.* The long DNA fragments cloned with the aid of cosmids can in turn be subcloned in the usual vectors suitable for sequencing and then sequenced, as is described e.g. by Sanger et al. (Proceedings of the National Academy of Sciences of the United States of America, 74:5463-5467, 1977) *I.B.R.*

[0030] The resulting DNA sequences can then be investigated with known algorithms or sequence analysis programs, such as e.g. that of Staden (*Nucleic Acids Research* 14, 217-232 (1986)) *I.B.R.*, that of Marck (*Nucleic Acids Research* 16, 1829-1836 (1988)) *I.B.R.* or the GCG program of Butler (*Methods of Biochemical Analysis* 39, 74-97 (1998)) *I.B.R.*

[0031] The new DNA sequence of *C. glutamicum* which codes for the ppgK gene and which, as SEQ ID No. 1, is a constituent of the present invention has been found. The amino acid sequence of the corresponding protein has furthermore been derived from the present DNA sequence by the methods described above. The resulting amino acid sequence of the ppgK gene product is shown in SEQ ID No. 2.

[0032] Coding DNA sequences which result from SEQ ID No. 1 by the degeneracy of the genetic code are also a constituent of the invention. In the same way, DNA

sequences which hybridize with SEQ ID No. 1 or parts of SEQ ID No. 1 are a constituent of the invention. Conservative amino acid exchanges, such as e.g. exchange of glycine for alanine or of aspartic acid for glutamic acid in proteins, are furthermore known among experts as "sense mutations" which do not lead to a fundamental change in the activity of the protein, i.e. are of neutral function. It is furthermore known that changes on the N and/or C terminus of a protein cannot substantially impair or can even stabilize the function thereof. Information in this context can be found by the expert, inter alia, in Ben-Bassat et al. (Journal of Bacteriology 169:751-757 (1987)) I.B.R., in O'Regan et al. (Gene 77:237-251 (1989)) I.B.R., in Sahin-Toth et al. (Protein Sciences 3:240-247 (1994)) I.B.R., in Hochuli et al. (Bio/Technology 6:1321-1325 (1988)) I.B.R. and in known textbooks of genetics and molecular biology. Amino acid sequences which result in a corresponding manner from SEQ ID No. 2 are also a constituent of the invention.

[0033] In the same way, DNA sequences which hybridize with SEQ ID No. 1 or parts of SEQ ID No. 1 are a constituent of the invention. Finally, DNA sequences which are prepared by the polymerase chain reaction (PCR) using primers which result from SEQ ID No. 1 are a constituent of the invention. Such oligonucleotides typically have a length of at least 15 nucleotides.

[0034] Instructions for identifying DNA sequences by means of hybridization can be found by the expert, inter alia, in the handbook "The DIG System Users Guide for Filter Hybridization" from Boehringer Mannheim GmbH (Mannheim, Germany, 1993) I.B.R. and in Liebl et al. (International Journal of Systematic Bacteriology (1991) 41: 255-260) I.B.R. The hybridization takes place under stringent conditions, that is to say only hybrids in which the probe and target sequence, i. e. the polynucleotides treated with the probe, are at least 70% identical are formed. It is known that the stringency of the hybridization, including

the washing steps, is influenced or determined by varying the buffer composition, the temperature and the salt concentration. The hybridization reaction is preferably carried out under a relatively low stringency compared with the washing steps (Hybaid Hybridisation Guide, Hybaid Limited, Teddington, UK, 1996 *I.B.R.*).

[0035] A 5x SSC buffer at a temperature of approx. 50°C - 68°C, for example, can be employed for the hybridization reaction. Probes can also hybridize here with polynucleotides which are less than 70% identical to the sequence of the probe. Such hybrids are less stable and are removed by washing under stringent conditions. This can be achieved, for example, by lowering the salt concentration to 2x SSC and optionally subsequently 0.5x SSC (The DIG System User's Guide for Filter Hybridisation, Boehringer Mannheim, Mannheim, Germany, 1995 *I.B.R.*) a temperature of approx. 50°C - 68°C being established. It is optionally possible to lower the salt concentration to 0.1x SSC. Polynucleotide fragments which are, for example, at least 70% or at least 80% or at least 90% to 95% identical to the sequence of the probe employed can be isolated by increasing the hybridization temperature stepwise from 50°C to 68°C in steps of approx. 1 - 2°C. Further instructions on hybridization are obtainable on the market in the form of so-called kits (e.g. DIG Easy Hyb from Roche Diagnostics GmbH, Mannheim, Germany, Catalogue No. 1603558).

[0036] Instructions for amplification of DNA sequences with the aid of the polymerase chain reaction (PCR) can be found by the expert, inter alia, in the handbook by Gait: Oligonucleotide Synthesis: A Practical Approach (IRL Press, Oxford, UK, 1984) *I.B.R.* and in Newton and Graham: PCR (Spektrum Akademischer Verlag, Heidelberg, Germany, 1994) *I.B.R.*

[0037] It has been found that coryneform bacteria produce amino acids in an improved manner after over-expression of the ppgK gene.

[0038] To achieve an over-expression, the number of copies of the corresponding genes can be increased, or the promoter and regulation region or the ribosome binding site upstream of the structural gene can be mutated. Expression cassettes which are incorporated upstream of the structural gene act in the same way. By inducible promoters, it is additionally possible to increase the expression in the course of fermentative amino acid production. The expression is likewise improved by measures to prolong the life of the m-RNA. Furthermore, the enzyme activity is also increased by preventing the degradation of the enzyme protein. The genes or gene constructs can either be present in plasmids with a varying number of copies, or can be integrated and amplified in the chromosome. Alternatively, an over-expression of the genes in question can furthermore be achieved by changing the composition of the media and the culture procedure.

[0039] Instructions in this context can be found by the expert, inter alia, in Martin et al. (Bio/Technology 5, 137-146 (1987)) I.B.R., in Guerrero et al. (Gene 138, 35-41 (1994)) I.B.R., Tsuchiya and Morinaga (Bio/Technology 6, 428-430 (1988)) I.B.R., in Eikmanns et al. (Gene 102, 93-98 (1991)) I.B.R., in European Patent Specification 0 472 869 I.B.R., in US Patent 4,601,893 I.B.R., in Schwarzer and Pühler (Bio/Technology 9, 84-87 (1991)) I.B.R., in Reinscheid et al. (Applied and Environmental Microbiology 60, 126-132 (1994)) I.B.R., in LaBarre et al. (Journal of Bacteriology 175, 1001-1007 (1993)) I.B.R., in Patent Application WO 96/15246 I.B.R., in Malumbres et al. (Gene 134, 15 - 24 (1993)) I.B.R., in Japanese Laid-Open Specification JP-A-10-229891 I.B.R., in Jensen and Hammer (Biotechnology and Bioengineering 58, 191-195 (1998)) I.B.R., in Makrides (Microbiological Reviews 60:512-538

(1996)) *I.B.R.* and in known textbooks of genetics and molecular biology.

[0040] By way of example, for enhancement the *ppgK* gene according to the invention was over-expressed with the aid of episomal plasmids. Suitable plasmids are those which are replicated in coryneform bacteria. Numerous known plasmid vectors, such as e.g. *pZ1* (Menkel et al., *Applied and Environmental Microbiology* (1989) 64: 549-554 *I.B.R.*), *pEKEx1* (Eikmanns et al., *Gene* 102:93-98 (1991) *I.B.R.*) or *pHS2-1* (Sonnen et al., *Gene* 107:69-74 (1991) *I.B.R.*) are based on the cryptic plasmids *pHM1519*, *pBL1* or *pGAl*. Other plasmid vectors, such as e.g. those based on *pCG4* (US-A 4,489,160 *I.B.R.*), or *pNG2* (Serwold-Davis et al., *FEMS Microbiology Letters* 66, 119-124 (1990) *I.B.R.*), or *pAG1* (US-A 5,158,891) *I.B.R.*, can be used in the same manner.

[0041] Plasmid vectors which are furthermore suitable are also those with the aid of which the process of gene amplification by integration into the chromosome can be used, as has been described, for example, by Reinscheid et al. (*Applied and Environmental Microbiology* 60, 126-132 (1994)) *I.B.R.* for duplication or amplification of the *hom-thrB* operon. In this method, the complete gene is cloned in a plasmid vector which can replicate in a host (typically *E. coli*), but not in *C. glutamicum*. Possible vectors are, for example, *pSUP301* (Simon et al., *Bio/Technology* 1, 784-791 (1983) *I.B.R.*), *pK18mob* or *pK19mob* (Schäfer et al., *Gene* 145, 69-73 (1994) *I.B.R.*), *pGEM-T* (Promega Corporation, Madison, WI, USA), *pCR2.1-TOPO* (Shuman (1994). *Journal of Biological Chemistry* 269:32678-84 *I.B.R.*; US-A 5,487,993 *I.B.R.*), *PCR@Blunt* (Invitrogen, Groningen, Holland; Bernard et al., *Journal of Molecular Biology*, 234: 534-541 (1993) *I.B.R.*), *pEM1* (Schrumpf et al, 1991, *Journal of Bacteriology* 173:4510-4516 *I.B.R.*) or *pBGS8* (Spratt et al., 1986, *Gene* 41: 337-342 *I.B.R.*). The plasmid vector which contains the gene to be amplified is then transferred into the desired strain of *C. glutamicum* by conjugation or

transformation. The method of conjugation is described, for example, by Schäfer et al. (Applied and Environmental Microbiology 60, 756-759 (1994)) *I.B.R. Methods for transformation* are described, for example, by Thierbach et al. (Applied Microbiology and Biotechnology 29, 356-362 (1988)) *I.B.R.*, Dunican and Shivan (Bio/Technology 7, 1067-1070 (1989)) *I.B.R.* and Tauch et al. (FEMS Microbiological Letters 123, 343-347 (1994)) *I.B.R.* After homologous recombination by means of a "cross over" event, the resulting strain contains at least two copies of the gene in question.

[0042] In addition, it may be advantageous for the production of L-amino acids to enhance, in particular over-express, one or more enzymes of the particular biosynthesis pathway, of glycolysis, of anaplerosis, of the citric acid cycle, of the pentose phosphate cycle, of amino acid export and optionally regulatory proteins, in addition to the ppgK gene.

[0043] Thus, for the preparation of L-amino acids, in addition to enhancement of the ppgK gene, one or more endogenous genes chosen from the group consisting of

- the dapA gene which codes for dihydrodipicolinate synthase (EP-B 0 197 335 *I.B.R.*),
- the gap gene which codes for glyceraldehyde 3-phosphate dehydrogenase (Eikmanns (1992), Journal of Bacteriology 174:6076-6086 *I.B.R.*),
- the tpi gene which codes for triose phosphate isomerase (Eikmanns (1992), Journal of Bacteriology 174:6076-6086 *I.B.R.*),
- the pgk gene which codes for 3-phosphoglycerate kinase (Eikmanns (1992), Journal of Bacteriology 174:6076-6086 *I.B.R.*),

- the zwf gene which codes for glucose 6-phosphate dehydrogenase (JP-A-09224661 *I.B.R.*),
- the pyc gene which codes for pyruvate carboxylase (DE-A-198 31 609 *I.B.R.*),
- the mqo gene which codes for malate-quinone oxidoreductase (Molenaar et al., European Journal of Biochemistry 254, 395-403 (1998) *I.B.R.*),
- the lysC gene which codes for a feed-back resistant aspartate kinase (Accession No.P26512),
- the lysE gene which codes for lysine export (DE-A-195 48 222 *I.B.R.*),
- the hom gene which codes for homoserine dehydrogenase (EP-A 0131171 *I.B.R.*),
- the ilvA gene which codes for threonine dehydratase (Möckel et al., Journal of Bacteriology (1992) 8065-8072) *I.B.R.*) or the ilvA(Fbr) allele which codes for a "feed back resistant" threonine dehydratase (Möckel et al., (1994) Molecular Microbiology 13: 833-842 *I.B.R.*),
- the ilvBN gene which codes for acetohydroxy-acid synthase (EP-B 0356739 *I.B.R.*),
- the ilvD gene which codes for dihydroxy-acid dehydratase (Sahm and Eggeling (1999) Applied and Environmental Microbiology 65: 1973-1979 *I.B.R.*),
- the zwal gene which codes for the Zwal protein (DE: 19959328.0 *I.B.R.*, DSM 13115),

can be enhanced, in particular over-expressed.

[0044] It may furthermore be advantageous for the production of L-amino acids, in addition to the enhancement of the ppgK gene, for one or more genes chosen from the group consisting of:

- the *pck* gene which codes for phosphoenol pyruvate carboxykinase (DE 199 50 409.1 *I.B.R.*; DSM 13047; EP-A-1 094 111 *I.B.R.*),
- the *pgi* gene which codes for glucose 6-phosphate isomerase (US 09/396,478 *I.B.R.*; DSM 12969),
- the *poxB* gene which codes for pyruvate oxidase (DE: 1995 1975.7 *I.B.R.*; DSM 13114),
- the *zwa2* gene which codes for the Zwa2 protein (DE: 19959327.2 *I.B.R.*, DSM 13113)

to be attenuated, in particular for the expression thereof to be reduced.

[0045] The term "attenuation" used in this context describes the reduction or switching off of the intracellular activity of one or more enzymes (proteins) in a microorganism that are coded by the corresponding DNA, by for example using a weak promoter or using a gene or allele that codes for a corresponding enzyme having a low activity or that inactivates the corresponding gene or enzyme (protein), and optionally combining these measures.

[0046] By such attenuation measures the activity or concentration of the corresponding protein is in general reduced to 0 to 75%, 0 to 50%, 0 to 25%, 0 to 10% or 0 to 5% of the activity or concentration of the wild type protein, or the activity or concentration of the protein in the starting microorganism.

[0047] In addition to over-expression of the *ppgK* gene it may furthermore be advantageous for the production of amino acids to eliminate undesirable side reactions (Nakayama: "Breeding of Amino Acid Producing Micro-organisms", in: Overproduction of Microbial Products, Krumphanzl, Sikyta, Vanek (eds.), Academic Press, London, UK, 1982 *I.B.R.*).

[0048] The invention also provides the microorganisms prepared according to the invention, and these can be cultured continuously or discontinuously in the batch process (batch culture) or in the fed batch (feed process) or repeated fed batch process (repetitive feed process) for the purpose of production of amino acids. A summary of known culture methods is described in the textbook by Chmiel (Bioprozesstechnik 1. Einführung in die Bioverfahrenstechnik [Bioprocess Technology 1. Introduction to Bioprocess Technology (Gustav Fischer Verlag, Stuttgart, 1991)) I.B.R. or in the textbook by Storhas (Bioreaktoren und periphere Einrichtungen [Bioreactors and Peripheral Equipment] (Vieweg Verlag, Braunschweig/ Wiesbaden, 1994)) I.B.R.

[0049] The culture medium to be used must meet the requirements of the particular strains in a suitable manner. Descriptions of culture media for various microorganisms are contained in the handbook "Manual of Methods for General Bacteriology" of the American Society for Bacteriology (Washington D.C., USA, 1981) I.B.R.

[0050] Sugars and carbohydrates, such as e.g. glucose, sucrose, lactose, fructose, maltose, molasses, starch and cellulose, oils and fats, such as e.g. soya oil, sunflower oil, groundnut oil and coconut fat, fatty acids, such as e.g. palmitic acid, stearic acid and linoleic acid, alcohols, such as e.g. glycerol and ethanol, and organic acids, such as e.g. acetic acid, can be used as the source of carbon. These substance can be used individually or as a mixture.

[0051] Organic nitrogen-containing compounds, such as peptones, yeast extract, meat extract, malt extract, corn steep liquor, soya bean flour and urea, or inorganic compounds, such as ammonium sulfate, ammonium chloride, ammonium phosphate, ammonium carbonate and ammonium

nitrate, can be used as the source of nitrogen. The sources of nitrogen can be used individually or as a mixture.

[0052] Phosphoric acid, potassium dihydrogen phosphate or dipotassium hydrogen phosphate or the corresponding sodium-containing salts can be used as the source of phosphorus. The culture medium must furthermore comprise salts of metals, such as e. g. magnesium sulfate or iron sulfate, which are necessary for growth. Finally, essential growth substances, such as amino acids and vitamins, can be employed in addition to the above-mentioned substances. Suitable precursors can moreover be added to the culture medium. The starting substances mentioned can be added to the culture in the form of a single batch, or can be fed in during the culture in a suitable manner.

[0053] Basic compounds, such as sodium hydroxide, potassium hydroxide, ammonia or aqueous ammonia, or acid compounds, such as phosphoric acid or sulfuric acid, can be employed in a suitable manner to control the pH of the culture. Antifoams, such as e.g. fatty acid polyglycol esters, can be employed to control the development of foam. Suitable substances having a selective action, such as e.g. antibiotics, can be added to the medium to maintain the stability of plasmids. To maintain aerobic conditions, oxygen or oxygen-containing gas mixtures, such as e.g. air, are introduced into the culture. The temperature of the culture is usually 20°C to 45°C, and preferably 25°C to 40°C. Culturing is continued until a maximum of the desired product has formed. This target is usually reached within 10 hours to 160 hours.

[0054] Methods for the determination of L-amino acids are known from the prior art. The analysis can thus be carried out, for example, as described by Spackman et al. (Analytical Chemistry, 30, (1958), 1190) I.B.R. by ion exchange chromatography with subsequent ninhydrin derivatation, or it can be carried out by reversed phase

HPLC, for example as described by Lindroth et al.
(Analytical Chemistry (1979) 51: 1167-1174) *I.B.R.*

[0055] The process according to the invention is used for fermentative preparation of amino acids.

[0056] The present invention is explained in more detail in the following with the aid of embodiment examples.

[0057] The isolation of plasmid DNA from *Escherichia coli* and all techniques of restriction, Klenow and alkaline phosphatase treatment were carried out by the method of Sambrook et al. (Molecular Cloning. A Laboratory Manual (1989) Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, USA) *I.B.R.* Methods for transformation of *Escherichia coli* are also described in this handbook.

[0058] The composition of the usual nutrient media, such as LB or TY medium, can also be found in the handbook by Sambrook et al.

[0059] Example 1

Preparation of a genomic cosmid gene library from *Corynebacterium glutamicum* ATCC 13032

[0060] Chromosomal DNA from *Corynebacterium glutamicum* ATCC 13032 was isolated as described by Tauch et al. (1995, Plasmid 33:168-179) *I.B.R.* and partly cleaved with the restriction enzyme *Sau3AI* (Amersham Pharmacia, Freiburg, Germany, Product Description *Sau3AI*, Code no. 27-0913-02). The DNA fragments were dephosphorylated with shrimp alkaline phosphatase (Roche Diagnostics GmbH, Mannheim, Germany, Product Description *SAP*, Code no. 1758250). The DNA of the cosmid vector *SuperCos1* (Wahl et al. (1987) Proceedings of the National Academy of Sciences USA 84:2160-2164 *I.B.R.*), obtained from Stratagene (La Jolla, USA, Product Description *SuperCos1* Cosmid Vector Kit, Code no. 251301) was cleaved with the restriction enzyme *XbaI*

(Amersham Pharmacia, Freiburg, Germany, Product Description XbaI, Code no. 27-0948-02) and likewise dephosphorylated with shrimp alkaline phosphatase.

[0061] The cosmid DNA was then cleaved with the restriction enzyme BamHI (Amersham Pharmacia, Freiburg, Germany, Product Description BamHI, Code no. 27-0868-04). The cosmid DNA treated in this manner was mixed with the treated ATCC13032 DNA and the batch was treated with T4 DNA ligase (Amersham Pharmacia, Freiburg, Germany, Product Description T4-DNA-Ligase, Code no.27-0870-04). The ligation mixture was then packed in phages with the aid of Gigapack II XL Packing Extracts (Stratagene, La Jolla, USA, Product Description Gigapack II XL Packing Extract, Code no. 200217).

[0062] For infection of the E. coli strain NM554 (Raleigh et al. 1988, Nucleic Acids Research 16:1563-1575 I.B.R.) the cells were taken up in 10 mM MgSO₄ and mixed with an aliquot of the phage suspension. The infection and titering of the cosmid library were carried out as described by Sambrook et al. (1989, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor) I.B.R., the cells being plated out on LB agar (Lennox, 1955, Virology, 1:190 I.B.R.) with 100 mg/l ampicillin. After incubation overnight at 37°C, recombinant individual clones were selected.

[0063] Example 2

Isolation and sequencing of the ppgK gene

[0064] The cosmid DNA of an individual colony was isolated with the Qiaprep Spin Miniprep Kit (Product No. 27106, Qiagen, Hilden, Germany) in accordance with the manufacturer's instructions and partly cleaved with the restriction enzyme Sau3AI (Amersham Pharmacia, Freiburg, Germany, Product Description Sau3AI, Product No. 27-0913-02). The DNA fragments were dephosphorylated with shrimp alkaline phosphatase (Roche Diagnostics GmbH, Mannheim,

Germany, Product Description SAP, Product No. 1758250). After separation by gel electrophoresis, the cosmid fragments in the size range of 1500 to 2000 bp were isolated with the QiaExII Gel Extraction Kit (Product No. 20021, Qiagen, Hilden, Germany).

[0065] The DNA of the sequencing vector pZero-1, obtained from Invitrogen (Groningen, Holland, Product Description Zero Background Cloning Kit, Product No. K2500-01), was cleaved with the restriction enzyme BamHI (Amersham Pharmacia, Freiburg, Germany, Product Description BamHI, Product No. 27-0868-04). The ligation of the cosmid fragments in the sequencing vector pZero-1 was carried out as described by Sambrook et al. (1989, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor) *I.B.R.*, the DNA mixture being incubated overnight with T4 ligase (Pharmacia Biotech, Freiburg, Germany). This ligation mixture was then electroporated (Tauch et al. 1994, FEMS Microbiol Letters, 123:343-7 *I.B.R.*) into the *E. coli* strain DH5 α MCR (Grant, 1990, Proceedings of the National Academy of Sciences U.S.A., 87:4645-4649) *I.B.R.* and plated out on LB agar (Lennox, 1955, Virology, 1:190 *I.B.R.*) with 50 mg/l zeocin.

[0066] The plasmid preparation of the recombinant clones was carried out with the Biorobot 9600 (Product No. 900200, Qiagen, Hilden, Germany). The sequencing was carried out by the dideoxy chain termination method of Sanger et al. (1977, Proceedings of the National Academy of Sciences U.S.A., 74:5463-5467) *I.B.R.* with modifications according to Zimmermann et al. (1990, Nucleic Acids Research, 18:1067) *I.B.R.* The "RR dRhodamin Terminator Cycle Sequencing Kit" from PE Applied Biosystems (Product No. 403044, Weiterstadt, Germany) was used. The separation by gel electrophoresis and analysis of the sequencing reaction were carried out in a "Rotiphoresis NF Acrylamide/Bisacrylamide" Gel (29:1) (Product No. A124.1, Roth, Karlsruhe, Germany) with the "ABI Prism 377"

sequencer from PE Applied Biosystems (Weiterstadt, Germany).

[0067] The raw sequence data obtained were then processed using the Staden program package (1986, Nucleic Acids Research, 14:217-231 *I.B.R.*) version 97-0. The individual sequences of the pZerol derivatives were assembled to a continuous contig. The computer-assisted coding region analysis was prepared using the program XNIP (Staden, 1986, Nucleic Acids Research, 14:217-231 *I.B.R.*). Further analyses can be carried out with the "BLAST search program" (Altschul et al., 1997, Nucleic Acids Research, 25:3389-3402 *I.B.R.*) against the non-redundant databank of the "National Center for Biotechnology Information" (NCBI, Bethesda, MD, USA) *I.B.R.*

[0068] The relative degree of substitution or mutation in the polynucleotide or amino acid sequence to produce a desired percentage of sequence identity can be established or determined by well-known methods of sequence analysis. These methods are disclosed and demonstrated in Bishop, et al. "DNA & Protein Sequence Analysis (A Practical Approach)", Oxford Univ. Press, Inc. (1997) *I.B.R.* and by Steinberg, Michael "Protein Structure Prediction" (A Practical Approach), Oxford Univ. Press, Inc. (1997) *I.B.R.*

[0069] The resulting nucleotide sequence is shown in SEQ ID No. 1. Analysis of the nucleotide sequence showed an open reading frame of 789 base pairs, which was called the ppgK gene. The ppgK gene codes for a protein of 262 amino acids.

[0070] This application claims priority to German Priority Document Application No. 100 47 403.9, filed on September 26, 2000. The above German Priority Document is hereby incorporated by reference in its entirety.